PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



(51) International Patent Classification 7:		NDER THE PATENT COOPERATION TREATY (PCT) (11) International Publication Number: WO 00/12043				
A61J 1/05, 1/14	A1					
		(43) International Publication Date: 9 March 2000 (09.03.00)				
(21) International Application Number: PCT/SE (22) International Filing Date: 24 August 1999 ((30) Priority Data: 9802938-2 1 September 1998 (01.09.98) (71) Applicant (for all designated States except US): AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE (72) Inventors; and (75) Inventors/Applicants (for US only): LUNDGRE [SE/SE]; Astra Hässle AB, S-431 83 Mölne SUNDGREN, Mats [SE/SE]; Astra Hässle AB, Mölndal (SE). (74) Agent: ASTRA AKTIEBOLAG; Intellectual Property S-151 85 Södertälje (SE).	(24.08.9 8) S ASTR E). EN, And dal (SE S-431	BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of				
(54) Title: IMPROVED STABILITY FOR INJECTION	SOLUT	ONS .				
(57) Abstract						
(67) 1250 401						
	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				
A primary package containing a low molecular we	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				
A primary package containing a low molecular we	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				
A primary package containing a low molecular we	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				
A primary package containing a low molecular we	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				
A primary package containing a low molecular we	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				
A primary package containing a low molecular we	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				
A primary package containing a low molecular we	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				
A primary package containing a low molecular we	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				
A primary package containing a low molecular we	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				
A primary package containing a low molecular we	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				
A primary package containing a low molecular we	eight pe	tide-based thrombin inhibitors which package is sealed with a rubbe				

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	Fl	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
ВЈ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Vict Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	Pl.	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

WO 00/12043 PCT/SE99/01440

IMPROVED STABILITY FOR INJECTION SOLUTIONS

Field of the invention

The present invention relates to solutions of low molecular weight thrombin inhibitors stored in primary packages containing rubber components, such as vials, bottles, cartridges and prefilled syringes. The invention also relates to the medical use of such stored thrombin inhibitor solutions.

10 Background of the invention

Solutions for parentheral use of pharmaceutically active substances are normally stored in primary packages such as, vials, bottles, cartridges or in prefilled syringes. The primary packages are sealed by a rubber stopper or plunger. A commonly used rubber material contains chlorobutyl. Solutions of low molecular weight thrombin inhibitors stored in vials, bottles, cartridges and prefilled syringes sealed by a stopper or plunger containing chlorobutyl rubber exhibits increased degradation, leading to shortened time of storage.

Disclosure of the invention

20

15

It has now surprisingly been found that by using rubber material containing bromobutyl instead of chlorobutyl, the stability of the low molecular weight thrombin inhibitors in solution can be considerably improved.

The present invention provides a primary package, such as a vial, a bottle, a cartridge or a prefilled syringe containing a solution of a low molecular weight thrombin inhibitor for parentheral injection, sealed by a rubber stopper or plunger containing bromobutyl rubber instead of chlorobutyl rubber.

WO 00/12043 PCT/SE99/01440

2

The present invention further provides a medical use of such thrombin inhibitor, or salts of such thrombin inhibitor, solutions kept in a primary package as mentioned above sealed by bromobutyl stoppers or plungers.

The present invention further provides an aqueous solution for parenteral administration comprising a low molecular weight peptide-based thrombin inhibitor or a salt thereof, having a pH in the range 3 to 8, preferably a pH about 5 and stored in a primary package, such as a vial, a bottle, a cartridge or a prefilled syringe, sealed by a rubber stopper or plunger containing bromobutyl.

10

Thrombin inhibitors referred to in this application are low molecular weight peptide-based thrombin inhibitors. The term "low molecular weight peptide-based thrombin inhibitors" will be well understood by one skilled in the art to include thrombin inhibitors with one to four peptide linkages, and/or with a molecular weight below 1000, and includes those described generically and, more preferably, specifically in the review paper by Claesson in Blood Coagul. Fibrin. (1994) 5, 411, as well as those disclosed in US Patent No. 4,346,078; International Patent Applications WO 97/23499, WO 97/02284, WO97/46577, WO 98/01422, WO 93/05069, WO93/11152, WO 95/23609, WO95/35309, WO 96/25426, WO 94/29336, WO WO 93/18060 and WO 95/01168; and European Patent Applications 623 596, 648 780, 468 231, 559 046, 641 779, 185 390, 526 877, 542 525, 195 212, 362 002, 364 344, 530 167, 293 881, 686 642, 669 317 and 601 459.

20

Preferred low molecular weight peptide-based thrombin inhibitors include those known collectively as the "gatrans". Particular gatrans which may be mentioned include HOOC-CH₂(R)Cha-Pic-Nag-H (known as inogatran; see International Patent Application WO 93/11152 and the list of abbreviations therein) and HOOC-CH₂-(R)Cgl-Aze-Pab-H (known as melagatran; see International Patent Application WO 94/29336 and the list of abbreviations therein).

The preferred low molecular weight peptide-based thrombin inhibitor to be kept in glass vials or syringes is selected from the group consisting of inogatran, (Glycine, N-[2-[2-[[[3-[(aminoimino-methyl)amino]propyl]amino]carbonyl]-1-piperidinyl]-1-(cyclohexylmethyl)-2-oxoethyl]-, [2R-[2S]]-), melagatran, (Glycine, N-[2-[2-[[[4-(aminoiminomethyl)phenyl]-methyl]amino]carbonyl]-1-azetidinyl]-1-cyclohexyl-2-oxoethyl]-, [2R-[2S]]-) and compound A, (Glycine, N-[1-cyclohexyl-2-[2-[[[4-[(hydroxyimino)aminomethyl]-phenyl]methyl]amino]carbonyl]-1-azetidinyl]-2-oxoethyl]-, ethyl ester, [S-(R*, S*)]-).

10

15

20

In one embodiment of the invention the thrombin inhibitor (preferably melagatran) solutions for parentheral injection is a water solution and are kept in primary packages such as vials, bottles, cartridges or prefilled syringes having a rubber stopper or plunger containing bromobutyl.

In another embodiment of the invention; the thrombin inhibitor for parentheral injection is in a water solution with an addition of hydroxy-propyl- β -cyclodextrin (HP β CD). The concentration of the thrombin inhibitor is in the range 0.001-100 mg/ml, preferably 2.5-20 mg/ml.

Working Example

Analytical technique

25

Liquid Chromatography (LC), for all analysis

The following equipment and parameters were used at the analysis of melagatran in solution.

Flowrate

1.0 ml/min

Wavelength

237 nm

Injection volume

20 µl

Analytical column

Waters Symmetry C8, 150 x 3.9 mm

Guard column

Waters Symmetry C8, 22 x 3.9 mm

Mobile phase

20 % (v/v) acetonitrile in phosphate buffer, pH 2.0 with 4.6 mM

octanesulphonic acid.

EVALUATION

10

Results in tables are presented as total degradation of melagatran. This means that all by-products are included and presented as area% of melagatran.

Example 1.

15

This example shows a comparison of melagatran in HP β CD-solution in prefilled syringes (1.0 ml) having rubber plungers containing bromobutyl and chlorobutyl, respectively. The syringes were stored at 4, 25 and 50 °C for up to 6 months.

The melagatran solution was in direct contact with the different rubber materials.

MANUFATURING OF SAMPLES

Melagatran, 2.5 mg/ml, in HPβCD water solution (40 % w/w), pH about 5

25 Batch HF 839-2601

Melagatran

442.1 mg

HPβCD

80.0 g

HCl, 1 M

qs

₃₀ NaOH, 1 M

qs

WO 00/12043 PCT/SE99/01440⁻

5

water for injection

to 200 g final weight (density 1.145 g/ml)

Melagatran was dissolved in water in a separate beaker and adjusted to pH 5.06. HP β CD powder was mixed with this solution together with water. The final solution was mixed with a magnetic stirrer until the substance was completely dissolved and pH was finally adjusted to 5.02, and the solution was filtrated with a 0.22 μ m sterile filter.

Melagatran, 10 mg/ml, in HPβCD water solution (40 % w/w), pH about 5

10 Batch HF 839-2602

20

Melagatran 1.77 mg

HPβCD 80.0 g

HCl, 1 M qs

NaOH, 1 M qs

water for injection to 200 g final weight (density 1.145 g/ml)

Melagatran was dissolved in water in a separate beaker and adjusted to pH 4.88. HP β CD powder was mixed with this solution together with water. The final solution was mixed with a magnetic stirrer until the substance was completely dissolved and pH was finally adjusted to 5.0, and the solution was filtrated with a 0.22 μ m sterile filter.

FILLING OF SYRINGES (1.0 ml)

25 Sample A1 (HF 839-2613) 10 mg/ml

0.5 ml of HF 839-2602 was filled in 1 ml HYPAK® syringes from Becton Dickinson with a black plunger material (PH 701/50 from The West Company) containing chlorobutyl rubber.

10

15

Sample B1 (HF 839-2614) 10 mg/ml

0.5 ml of HF 839-2602 was filled in 1 ml HYPAK® syringes from Becton Dickinson with a grey plunger material (PH 4416/50 from The West Company) containing bromobutyl rubber.

Sample C1 (HF 839-2615) 2.5 mg/ml

0.5 ml of HF 839-2601 was filled in 1 ml HYPAK® syringes from Becton Dickinson with, a grey plunger material (PH 4416/50 from The West Company) containing bromobutyl rubber.

Sample D1 (HF 839-2616) 10 mg/ml

0.5 ml of HF 839-2602 was filled in 1 ml HYPAK® syringes from Becton Dickinson with a black plunger material (PH 701/50 from The West Company) containing chlorobutyl rubber.

RESULTS OF STABILITY STUDIES

Sample A1 (HF 839-2613) 10 mg/ml - Chlorobutyl rubber

Storage time (months)	pН	Temperature (°C)	Total degradation (area % of melagatran)
0	5.2	-	1.2
1	5.2	4	1.0
1	5.3	50	7.4
3	5.1	4	1.2
3	5.1	25	4.5
3	5.2	50	14.9
6	5.1	4	1.2
6	5.1	25	3.7

Sample B1 (HF 839-2614) 10 mg/ml - Bromobutyl rubber

Sample 21 (111		T	
Storage time	pН	Temperature	Total degradation
(months)		(°C)	(area % of melagatran)
0	5.2		1.1
1	5.2	4	1.0
1	5.2	50	6.41
3	5.1	4	1.2
3	5.1	25	2.4
3	5.2	50	12.8
6	5.1	4	1.1
6	5.1	25	3.1

Sample C1 (HF 839-2615) 2.5 mg/ml - Bromobutyl rubber

Storage time	pН	Temperature	Total degradation
(months)		(°C)	(area % of melagatran)
0	5.3		1.2
1	5.4	4	1.1
1	5.3	50	7.2
3	5.3	4	1.3
3	5.3	25	3.9
3	5.2	50	14.2
6	5.2	4	1.2
6	5.2	25	5.7

Sample D1 (HF 839-2616) 10 mg/ml - Chlorobutyl rubber

Sample D1 (HF 839-261)	6) 10 mg/m - Chlores	1	
Storage time	pН	Temperature	Total degradation
(months)		(°C)	(area % of melagatran)
0	5.3	-	1.2
1	5.4	4	1.2
1	5.3	50	8.6
3	5.3	4	1.2:
3	5.3	25	3.1
3	5.2	50	17.4
6	5.2	4	1.4
6	5.2	25	9.9

Conclusion

Rubber plungers containing chlorobutyl result in a more pronounced degradation compared to rubber plungers containing bromobutyl. This is true for high concentrations as well as low concentrations of melagatran in aqueous solutions.

The most pronounced difference was seen between plungers of chlorobutyl rubber and bromobutyl rubber when the dose of melagatran in aqueous solution was as low as 2.5 mg/ml.

Example 2.

This example is a comparison of melagatran in a water solution of HPβCD and melagatran in a water solution of NaCl. Both solutions are in direct contact with rubber plungers containing bromobutyl.

3 plungers of the quality FM 257 (from Helvoet Pharma N.V.) were placed in each 3 ml glass vial together with 1 ml solution of melagatran (NaCl water solution and HPβCD water solution, respectively). Reference samples, that is melagatran in NaCl water solution and in HPβCD water solution having no contact with plunger material. The reference samples were treated in the same way as the other samples. The vials were stored at 50 °C for up to 3 months.

Compared to the study of Example 1 the ratio between solution exposed plunger surface and the quantity of melagatran solution is 16 times higher.

10

MANUFATURING OF SAMPLES

Melagatran, 7.5 mg/ml, in HP β CD water solution (40 % w/w), pH about 5.

Batch HF 839-2679

15 Melagatran 928.8 mg
HPβCD 55.0 g

HCl, 1 M qs

NaOH, 1 M qs

water for injection 137.4 g (density 1.145 g/ml)

20

Melagatran and HP β CD were dissolved in water and adjusted to pH 4.96. The final solution was diluted with water to final weight and sterile filtrated with 0.45 μ m filter.

qs

Melagatran, 7.5 mg/ml, in NaCl water solution, pH about 5.

Batch HF 839-2680

Melagatran 1315.5g

NaCl 1.441 g

HCl, 1 M

NaOH, 1 M

qs

water for injection

to 170 (density 1.0 g/ml)

Melagatran and NaCl were dissolved in water and adjusted to pH 5.03. The final solution was diluted with water to final weight and sterile filtrated with 0.22 μ m filter.

FILLING OF VIALS

20

Sample A2 (HF 839-2682) 7.5 mg/ml in NaCl

1.0 ml of HF 839-2680 was filled in 3 ml vials together with 3 black unsiliconized plungers (FM 257 from Helvoet Pharma N.V.) containing bromobutyl rubber.

Sample B2 (HF 839-2683) 7.5 mg/ml in NaCl

1.0 ml of HF 839-2680 was filled in 3 ml vials together with 3 black siliconized plungers
(FM 257 from Helvoet Pharma N.V.) containing bromobutyl rubber.

Sample C2 (HF 839-2684) 7.5 mg/ml in NaCl

1.0 ml of HF 839-2680 was filled in 3 ml vials together with 3 grey siliconized plungers (FM 257 from Helvoet Pharma N.V.) containing bromobutyl rubber.

Sample D2 (HF 839-2688) 7.5 mg/ml in NaCl

1.0 ml of HF 839-2680 was filled in 3 ml vials (Reference).

Sample E2 (HF 839-2689) 7.5 mg/ml in HPβCD

1.0 ml of HF 839-2679 was filled in 3 ml vials together with 3 black unsiliconized plungers (FM 257 from Helvoet Pharma N.V.) containing bromobutyl rubber.

Sample F2 (HF 839-2690) 7.5 mg/ml in HPβCD

1.0 ml of HF 839-2679 was filled in 3 ml vials together with 3 black siliconized plungers (FM 257 from Helvoet Pharma N.V.) containing bromobutyl rubber.

Sample G2 (HF 839-2691) 7.5 mg/ml in HP β CD

1.0 ml of HF 839-2679 was filled in 3 ml vials together with 3 grey siliconized plungers (FM 257 from Helvoet Pharma N.V.) containing bromobutyl rubber.

Sample H2 (HF 839-2695) 7.5 mg/ml in HP β CD

1.0 ml of HF 839-2679 was filled in 3 ml vials (Reference).

RESULTS OF STABILITY STUDIES

10

5

Sample A2 (HF 839-2682) 7.5 mg/ml in NaCl - Bromobutyl rubber

		Total degradation
ргı	_	(area % of melagatran)
· · · · · · · · · · · · · · · · · · ·	(0)	
5.9	50	4.2
6.0	50	9.3
	рН	(°C) 5.9 50

Sample B2 (HF 839-2683) 7.5 mg/ml in NaCl - Bromobutyl rubber

Sample B2 (Hr 839-20	83) 7.5 mg/mi iii 14aCi		
Storage time	pН	Temperature	Total degradation
(months)		(°C)	(area % of melagatran)
1	5.8	50	4.0
3	6.0	50	8.7
,			

15

Sample C2 (HF 839-2684) 7.5 mg/ml in NaCl - Bromobutyl rubber

Sample C2 (HF 839-2684	7.5 mg/ml in NaCl	- Bromobutyi rubbei	
Storage time	pН	Temperature	Total degradation
(months)		(°C)	(area % of melagatran)
1	5.8	50	3.7
2	5.8	50	7.9
3	5.0		

WO 00/12043

5

Sample D2 (HF 839-2688) 7.5 mg/ml in NaCl - Reference

Storage time	pН	Temperature	Total degradation
(months)		(°C)	(area % of melagatran)
1	5.2	4	1.4
3	5.3	4	1.4
1	5.4	50	3.4
3	5.6	50	6.8

Sample E2 (HF 839-2689) 7.5 mg/ml in HPBCD - Bromobutyl rubber

Storage time	pH	Temperature	Total degradation
(months)		(°C)	(area % melagatran)
1	5.5	50	5.5
3	5.6	50	11.3

Sample F2 (HF 839-2690) 7.5 mg/ml in HPβCD - Bromobutyl rubber

Storage time	pН	Temperature	Total degradation
(months)		(°C)	(area % of melagatran)
1	5.4	50	5.4
3	5.5	50	11.3

Sample G2 (HF 839-2691) 7.5 mg/ml in HPBCD - Bromobutyl rubber

Sample GZ (III 057-207	1) 1.5 mg/m in 222 p = 2		
Storage time	pН	Temperature	Total degradation
(months)		(°C)	(area % of melagatran)
1	5.4	50	5.4
3	5.5	50	10.3

Sample H2 (HF 839-2695) 7.5 mg/ml in HPBCD - Reference

Storage time	рН	Temperature	Total degradation
(months)		(°C)	(area % of melagatran)
1	5.2	4	1.5
3	5.3	4	1.7
1	5.3	. 50	5.7
3	5.4	50	10.7

Conclusion

5

10

Melagatran in a water solution of NaCl exhibits a somewhat lower degradation compared to melagatran in a water solution of HPβCD. This is true both for solutions in contact with plunger material (FM 257 bromobutyl) 8%* compared to 11%*, and solutions in absence of plunger material (reference) 7%* compared to 11%*.

*; is total degradation in area% of melagatran

Example 3.

This example shows a comparison of different kinds of stopper and plunger materials containing either bromobutyl rubber or chlorobutyl rubber in contact with a melagatran solution (NaCl, pH 5). Melagatran solution was filled in glass vials (3 ml) together with stoppers and plungers of different brands. 5 different rubber materials were used in the study. There were 3 different bromobutyl and 2 different chlorobutyl rubbers. As reference, NaCl water solution of melagatran was stored without any contact with stopper or plunger material.

The ratio between exposed plunger or stopper surface and melagatran in water solution is higher than in Example 1. A calculation has been made of exposed area of each tested plunger or stopper material. In the study the area ratio is 10-15 times higher compared to the area represented in Example 1. The vials were studied up to 19 days at a temperature of 50°C.

MANUFACTURING OF SAMPLES

Melagatran, 5 mg/ml, in isotonic NaCl solution, pH about 5.

Batch HF 839-2719

5

10

20

Melagatran 10.0 mg

NaCl 17.6 g

HCl, 1 M qs

NaOH, 1 M qs

water for injection To 2000 g final weight (density 1.0 g/ml)

Melagatran and NaCl were dissolved in water and pH adjusted to 4.95 The solution was diluted to final weight with water.

FILLING OF VIALS

The total contact surface between the rubber material and the solution was enhanced in different ways and different extent. One way was by putting pieces of vial stopper material into each vial. For sample A3, the stopper material was divided into eight equal parts, and two parts in each vial (total of 2/8). Another way to enhance the contact surface was to put 2-3 plungers in each vial. For sample E3, three plungers were put in each vial. In samples A3 to F3, the contact surface was increased of 10-15 times compared to the normal contact surface between plunger and solution in a 1 ml syringe (used in Example 1).

10

20

Sample A3 (HF 839-2727) 5 mg/ml in NaCl

1.5 ml of HF 839-2719 was filled in a 3 ml vial together with two 1/8 parts of a 10 ml vial stopper (FM 50 from Helvoet Pharma N.V.) containing chlorobutyl rubber.

Sample B3 (HF 839-2728) 5 mg/ml in NaCl

1.5 ml of HF 839-2719 was filled in 3 ml vial together with 2 grey plungers (PH 4023/50 from The West Company) containing bromobutyl rubber.

Sample C3 (HF 839-2729) 5 mg/ml in NaCl

1.5 ml of HF 839-2719 was filled in 3 ml vial together with 2 black plungers (PH 701/50 from The West Company) containing chlorobutyl rubber.

Sample D3 (HF 839-2730) 5 mg/ml in NaCl

1.5 ml of HF 839-2719 was filled in 3 ml vial together with 2 grey plungers (W 4416/50 from The West Company) containing bromobutyl rubber.

Sample E3 (HF 839-2731) 5 mg/ml in NaCl

1.5 ml of HF 839-2719 was filled in 3 ml vial together with 3 black plungers (FM 257 from Helvoet Pharma N.V.) containing bromobutyl rubber.

Sample F3 (HF 839-2732) 5 mg/ml in NaCl

1.5 ml of HF 839-2719 was filled in 3 ml vial (Reference).

RESULTS OF STABILITY STUDIES

Sample A3 (HF 839-2727) 5 mg/ml in NaCl - Chlorobutyl rubber

Sample A5 (III 05) 272	, , , , , , , , , , , , , , , , , , , ,			
Storage time	pН	Temperature	Total degradation	
(days)		(°C)	(area % of melagatran)	
11	~5.0	50	8.0	
19	~5.0	50	11.8	

5 Sample B3 (HF 839-2728) 5 mg/ml in NaCl - Bromobutyl rubber

Storage time	pН	Temperature	Total degradation
(days)		(°C)	(area % of melagatran)
11	~5.0	50	0.9
19	~5.0	50	1.4

Sample C3 (HF 839-2729) 5 mg/ml in NaCl - Chlorobutyl rubber

Dampie Co (XX 00) 1715	<u> </u>		T	
Storage time	pН	Temperature	Total degradation	
(days)		(°C)	(area % of melagatran)	
11	~5.0	50	1.5	
19	~5.0	50	2.4	

Sample D3 (HF 839-2730) 5 mg/ml in NaCl - Bromobutyl rubber

Sumple 20 (222 00) = 101				
Storage time	pН	Temperature	Total degradation	
(days)		(°C)	(area % of melagatran)	
11	~5.0	50	1.3	
19	~5.0	50	1.6	

Sample E3 (HF 839-2731) 5 mg/ml in NaCl - Bromobutyl rubber

Storage time	pН	Temperature	Total degradation
(days)		(°C)	(area % of melagatran)
11	~5.0	50	1.2
19	~5.0	50	1.4

Sample F3 (HF 839-2732) 5 mg/ml in NaCl - Reference

Storage time	pН	Temperature	Total degradation
(days)		(°C)	(area % of melagatran)
11	~5.0	50	0.6
19	~5.0	50	1.0

Conclusion

5

10

All three bromobutyl rubber materials demonstrate lower melagatran degradation compared to the two chlorobutyl rubber materials.

Summary conclusion

It is shown in Example 1 that, for water solutions containing melagatran stored in HYPAK® syringes (from Becton Dickinson), improved stability is demonstrated using plungers containing bromobutyl rubber compared to the corresponding plungers containing chlorobutyl rubber.

It is shown in Example 2 that, for water solutions of melagatran stored in glass vials, improved stability is demonstrated using a NaCl water solution compared to a HPβCD water solution. This is true for melagatran in solution with and without contact of plungers containing bromobutyl rubber.

WO 00/12043 PCT/SE99/01440

18

It is shown in Example 3 that for melagatran in a NaCl water solution, improved stability is demonstrated using rubber materials containing bromobutyl compared to rubber materials containing chlorobutyl.

CLAIMS

- 1. A primary package containing an aqueous solution for parenteral administration comprising a low molecular weight peptide-based thrombin inhibitor or a salt thereof, having a pH in the range 3 to 8, the primary package being sealed with a rubber stopper or plunger containing bromobutyl rubber.
 - 2. A primary package according to claim 1, wherein the primary package is a vial.

10

- 3. A primary package according to claim 1, wherein the primary package is a bottle.
- 4. A primary package according to claim 1, wherein the primary package is a cartridge.
- 5. A primary package according to claim 1, wherein the primary package is a prefilled syringe.
 - 6. A primary package according to any of the preceding claims, wherein the solution is a NaCl solution.

20

- 7. A primary package according to any of the preceding claims, wherein the solution also comprises hydroxy-propyl-β-cyclodextrin.
- 8. A primary package according to any of the preceding claims, wherein the concentration
 of the thrombin inhibitor in the solution is in the range 0.001-100 mg/ml, preferably 2.5-20 mg/ml.
 - 9. A primary package according to any of the preceding claims, wherein the pH of the solution is in the range 3-8.

PCT/SE99/01440

5

20

- 10. A primary package according to claim 9, wherein the pH of the solution is about 5.
- 11. A primary package according to any of the preceding claims, wherein the thrombin inhibitor is melagatran.
- 12. A primary package according to any of the preceding claims, wherein the thrombin inhibitor in the solution is inogatran.
- 13. A primary package according to any of the preceding claims, wherein the thrombin inhibitor in the solution is compound A.
 - 14. A primary package according to any of the preceding claims, wherein the bromobutyl rubber material consists of, or correspond to, the quality PH 4023/53.
- 15. A primary package according to any of claims 1-13, wherein the bromobutyl rubber material consists of, or correspond to, the quality W 4416/50.
 - 16. A primary package according to any of claims 1-13, wherein the bromobutyl rubber material consists of, or correspond to, the quality FM 257.
 - 17. Use of a rubber stopper or plunger containing bromobutyl rubber for sealing a primary package, such as a vial, a bottle, a cartridge or a prefilled syringe, containing a low molecular weight peptide-based thrombin inhibitor in an aqueous solution.
- 18. A process for the manufacture of a primary package according to claim 1 comprising the steps of dissolving a low molecular weight peptide-based thrombin inhibitor in an aqueous solution, adjusting the pH of the solution to be in the range 3 to 8, optionally adding a cyclodextrin substance, sterile filtering the solution and filling it on a primary package which is then sealed with a rubber stopper or plunger containing bromobutyl rubber.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 99/01440

		1	
A. CLASS	IFICATION OF SUBJECT MATTER		
IPC7: A	61J 1/05, A61J 1/14 International Patent Classification (IPC) or to both nation		
		onal classification and IPC	
	S SEARCHED commentation searched (classification system followed by commentation searched (classification system followed by comments of the c	Passification symbols)	
		nasanicadon symbolog	-
IPC7: A	ion searched other than minimum documentation to the e	when that each documents are included	in the fields searched
	TI, NO classes as above	Atent that such documents are also docu	
Electronic d	ata base consulted during the international search (name o	of data base and, where practicable, searc	ch terms used)
c. Docu	MENTS CONSIDERED TO BE RELEVANT		1
Category*	Citation of document, with indication, where appr	opriate, of the relevant passages	Relevant to claim No.
X	WO 9633216 A1 (PHARMACIA AB), 24 (24.10.96), page 6, line 17 -	October 1996 · line 19	1-18
,			
X	EP 0390244 A1 (DUPHAR INTERNATION 3 October 1990 (03.10.90), pa line 48 - line 51; page 3, li page 4, Table A	age 2,	1-18
A	WO 9739770 A1 (ASTRA AKTIEBOLAG) (30.10.97)	, 30 October 1997	1-18
Furt	her documents are listed in the continuation of Box	C. See patent family ann	ex.
"A" docum to be "E" erlier	al categories of cited documents: nent defining the general state of the art which is not considered of particular relevance document but published on or after the international filing date nent which may throw doubts on priority claim(s) or which is to establish the publication date of another citation or other	"T" later document published after the i date and not in conflict with the apithe principle or theory underlying the document of particular relevance: the considered novel or cannot be consisted when the document is taken also	plication but cited to understand he invention he claimed invention cannot be idered to involve an inventive one
specia "O" docur mean "P" docur	al reason (as specified) nent referring to an oral disclosure, use, exhibition or other s nent published prior to the international filing date but later than	"Y" document of particular relevance: it considered to involve an inventive: combined with one or more other sbeing obvious to a person skilled in "&" document member of the same pate	step when the document is uch documents, such combination the art
i	he actual completion of the international search	Date of mailing of the international	
27 De	cember 1999	Authorized officer	
Swedist Box 505	nd mailing address of the ISA/ n Patent Office 15, S-102 42 STOCKHOLM	Authorized officer Nebil Gecer/Els	n
	e No. + 46 8 666 02 86 /ISA/210 (second sheet) (July 1992)	Telephone No. + 46 8 782 25 0	<u> </u>

INTERNATIONAL SEARCH REPORT Information on patent family members

02/12/99

International application No. PCT/SE 99/01440

	atent document in search repor		Publication date		Patent family member(s)		Publication date
10	9633216	A1	24/10/96	AU	5412996 0871959		07/11/96 21/10/98
				EP IL		Ď	00/00/00
				JP	10510925	Ť	20/10/98
				SE	9501472	Ď	00/00/00
				US	5862196		19/01/99
EP	0390244	A1	03/10/90	SE	0390244	T3	
Li	0330211	•••	00, 20, 00	ĀT	84205	T	15/01/93
				CA	2012919	A	28/09/90
				DK	390244	T	08/02/93
				IL		A	07/10/94
				JP		A	20/11/90
				NL	0000		16/10/90
				US	5383864	A 	24/01/95
WO	9739770	A1	30/10/97	AU	2719597	A	12/11/97
				AU	7716496	A	19/06/97
				EP	0864173		16/09/98
				SE	9601556	D	00/00/00